

ON THE q -EXTENSIONS OF THE BERNOULLI AND EULER NUMBERS, RELATED IDENTITIES AND LERCH ZETA FUNCTION

TAEKYUN KIM, YOUNG-HEE KIM, AND KYUNG-WON HWANG

Abstract Recently, λ -Bernoulli and λ -Euler numbers are studied in [5, 10]. The purpose of this paper is to present a systematic study of some families of the q -extensions of the λ -Bernoulli and the λ -Euler numbers by using the bosonic p -adic q -integral and the fermionic p -adic q -integral. The investigation of these λ - q -Bernoulli and λ - q -Euler numbers leads to interesting identities related to these objects. The results of the present paper cover earlier results concerning q -Bernoulli and q -Euler numbers. By using derivative operator to the generating functions of λ - q -Bernoulli and λ - q -Euler numbers, we give the q -extensions of Lerch zeta function.

2000 Mathematics Subject Classification : 11S80, 11B68?

Key words and phrases : λ -Bernoulli numbers, λ -Euler numbers, p -adic q -integral, Lerch zeta function

1. INTRODUCTION, DEFINITIONS AND NOTATIONS

Throughout this paper, the symbols $\mathbb{Z}_p, \mathbb{Q}_p, \mathbb{C}$ and $\mathbb{C}_p$ denote the ring of p -adic rational integers, the field of p -adic rational numbers, the complex number field and the completion of algebraic closure of $\mathbb{Q}_p$, respectively. Let $\mathbb{N}$ be the set of natural numbers.

The symbol q can be treated as a complex number, $q \in \mathbb{C}$, or as a p -adic number, $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, then we always assume that $|q| < 1$. If $q \in \mathbb{C}_p$, then we usually assume that $|1 - q|_p < 1$. Here $|\cdot|_p$ stands for the p -adic absolute value in $\mathbb{C}_p$ with $|p|_p = \frac{1}{p}$. The q -basic natural numbers are defined by $[n]_q = \frac{1-q^{n+1}}{1-q} = 1 + q + q^2 + \cdots + q^n$ ($n \in \mathbb{N}$) and $[n]_{-q} = \frac{1-(-q)^{n+1}}{1+q}$. In this paper, we use the notation

$$[x]_q = \frac{1-q^{x+1}}{1-q} \quad \text{and} \quad [x]_{-q} = \frac{1-(-q)^{x+1}}{1+q}, \quad \text{see [1-19].}$$

Hence $\lim_{q \rightarrow 1} [x]_q = x$ for any x with $|x|_p \leq 1$ in the present p -adic case.

For $x \in \mathbb{Z}_p$, we say that g is a uniformly differentiable function at a point $a \in \mathbb{Z}_p$, and write $g \in UD(\mathbb{Z}_p)$, the set of uniformly differentiable function, if the difference quotients

$$F_g(x, y) = \frac{g(y) - g(x)}{y - x}$$

have a limit $l = g'(a)$ as $(x, y) \rightarrow (a, a)$. For $f \in UD(\mathbb{Z}_p)$, the q -deformed bosonic p -adic integral is defined as

$$(1) \quad I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} f(x) \frac{q^x}{[p^N]_q}, \quad \text{see [1-19],}$$

1

and the q -deformed fermionic p -adic integral is defined by

$$I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} f(x) \frac{(-q)^x}{[p^N]_{-q}}, \quad (\text{see [1-19]}).$$

For $n \in \mathbb{N}$, let $f_n(x) = f(x+n)$. Then

$$(2) \quad q^n I_{-q}(f_n) = (-1)^n I_{-q}(f) + [2]_q \sum_{l=0}^{n-1} (-1)^{n-1-l} q^l f(l).$$

The classical Bernoulli polynomials $B_n(x)$ and the Euler polynomials $E_n(x)$ are defined as

$$(3) \quad \frac{t}{e^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!} \quad \text{and} \quad \frac{2e^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}.$$

The Bernoulli numbers B_n and the Euler numbers E_n are defined as $B_n = B_n(0)$ and $E_n = E_n(0)$, (see [1-19]).

From (1), we note that

$$(4) \quad qI_q(f_1) = I_q(f) + (q-1)f(0) + \frac{q-1}{\log q} f'(0),$$

for $f_1(x) = f(x+1)$. By (4), we see that $I_1(f_1) = I_1(f) + f'(0)$, (see [7]).

Let u be algebraic in $\mathbb{C}_p$ (or $\mathbb{C}$). Then the Frobenius-Euler polynomials are defined as

$$(5) \quad \frac{1-u}{e^t - u} e^{xt} = \sum_{n=0}^{\infty} H_n(u, x) \frac{t^n}{n!}, \quad (\text{see [5]}).$$

In case $x = 0$, $H_n(u, 0) = H_n(u)$, which are called the Frobenius Euler numbers.

Let C_{p^n} be the cyclic group consisting of all p^n -th roots of unity in $\mathbb{C}_p$ for any $n \geq 0$ and T_p be the direct limit of C_{p^n} with respect to the natural morphisms, hence T_p is the union of all C_{p^n} with discrete topology.

For $\lambda \in T_p$ with $\lambda \neq 1$, if we use (4), then we have

$$(6) \quad \int_{\mathbb{Z}_p} e^{tx} \lambda^x d\mu_1(x) = \frac{t}{\lambda e^t - 1}.$$

From (6), the λ -Bernoulli numbers are defined as

$$(7) \quad \frac{t}{\lambda e^t - 1} = e^{B(\lambda)t} = \sum_{n=0}^{\infty} B_n(\lambda) \frac{t^n}{n!}, \quad (\text{see [5]})$$

with the usual convention of replacing $B^i(\lambda)$ by $B_i(\lambda)$. Thus, $B_k(\lambda)$ can be determined inductively by

$$(8) \quad \lambda(B(\lambda) + 1)^k - B_k(\lambda) = \begin{cases} 1, & \text{if } k = 1, \\ 0, & \text{if } k > 1, \end{cases} \quad (\text{see [5]}).$$

By the definition of the Frobenius-Euler numbers, we see that

$$(9) \quad \frac{t}{\lambda e^t - 1} = \sum_{m=0}^{\infty} \frac{1}{(m+1)!} \cdot \frac{(m+1)H_m(\lambda^{-1})}{\lambda - 1} t^{m+1}, \quad (\text{see [7]}).$$

For $m \geq 1$ and $\lambda \neq 1$, we have

$$(10) \quad B_m(\lambda) = \int_{\mathbb{Z}_p} x^m \lambda^x d\mu_1(x) = \frac{m}{\lambda - 1} H_{m-1}(\lambda^{-1}), \quad (\text{see [5]}).$$

We can also easily see that $\int_{\mathbb{Z}_p} \lambda^x d\mu_1(x) = 0$ and

$$e^t x = \lim_{m \rightarrow \infty} \sum_{\lambda \in C_{p^m}} \frac{t \lambda^x}{\lambda e^t - 1} = \sum_{n=0}^{\infty} \frac{t^n}{n!} \lim_{m \rightarrow \infty} \sum_{\lambda \in C_{p^m}} \int_{\mathbb{Z}_p} x^m \lambda^x d\mu_1(x) \lambda^x.$$

Consequently, we have

$$\begin{aligned} x^n &= B_n(1) + \sum_{\substack{\lambda \in T_p \\ \lambda \neq 1}} \frac{1}{\lambda - 1} H_{n-1}(\lambda^{-1}) \lambda^x \\ &= B_n(1) + \sum_{\substack{\lambda \in T_p \\ \lambda \neq 1}} \frac{B_n(\lambda)}{n} \lambda^x, \quad (\text{see [5]}). \end{aligned}$$

From (6) and (8), we note that

$$B_0(\lambda) = 0, B_1(\lambda) = \frac{1}{\lambda - 1}, B_2(\lambda) = -\frac{2\lambda}{(\lambda - 1)^2}, \dots$$

The Genocchi numbers are defined by the generating function

$$\frac{2t}{e^t + 1} = \sum_{n=0}^{\infty} G_n \frac{t^n}{n!}.$$

These numbers satisfy the relation $G_0 = 0, G_1 = 1, G_3 = G_5 = \dots = G_{2k+1} = 0$, and the even coefficients are $G_n = 2(1 - 2^n)B_n$.

For $\lambda \in \mathbb{C}_p$ with $|\lambda| < 1$, by (2), we have

$$(11) \quad \int_{\mathbb{Z}_p} \lambda^x e^{xt} d\mu_{-1}(x) = \frac{2}{\lambda e^t + 1}.$$

By (11), we define the λ -Euler numbers as follows :

$$(12) \quad \frac{2}{\lambda e^t + 1} = \sum_{n=0}^{\infty} \frac{E_n(\lambda)}{n!} t^n, \quad (\text{see [7, 9, 10]}).$$

Note that $E_n(\lambda) = \frac{2}{\lambda + 1} H_n(-\lambda^{-1})$.

From (12), we can easily derive

$$(13) \quad \int_{\mathbb{Z}_p} x^n \lambda^x d\mu_{-1}(x) = E_n(\lambda) = \frac{2}{\lambda + 1} H_n(-\lambda^{-1}).$$

The λ -Genocchi numbers are also defined as

$$t \int_{\mathbb{Z}_p} x^n \lambda^x d\mu_{-1}(x) = \frac{2t}{\lambda e^t + 1} = \sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!}.$$

Thus, we have $G_0(\lambda) = 0, G_1(\lambda) = \frac{2}{\lambda + 1}, \dots, E_n(\lambda) = \frac{G_{n+1}(\lambda)}{n+1}$.

In this paper, we study the q -extension of λ -Bernoulli number and λ -Euler numbers related to Lerch zeta function. The purpose of this paper is to present a systematic study of some families of the q -extension of the λ -Bernoulli and λ -Euler numbers by using the bosonic p -adic q -integral and the fermionic p -adic q -integral.

The investigation of these λ - q -Bernoulli and λ - q -Euler numbers leads to interesting identities related to these objects. The results of the present paper cover earlier results concerning q -Bernoulli and q -Euler numbers. By using derivative operator to the generating functions of λ - q -Bernoulli and λ - q -Euler numbers, we can give the q -extension of Lerch zeta function.

2. q -EXTENSION OF λ -BERNOULLI NUMBERS AND POLYNOMIALS

For $\lambda \in T_p$, let us consider the q -extension of λ -Bernoulli numbers as follows.

$$(14) \quad \beta_{k,q}(\lambda) = \int_{\mathbb{Z}_p} \lambda^x [x]_q^k d\mu_q(x).$$

From (14), we note that

$$\begin{aligned} \beta_{k,q}(\lambda) &= \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} \lambda^x [x]_q^k q^x \\ &= \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} (\lambda q)^x \left(\sum_{l=0}^k \binom{k}{l} (-1)^l q^{lx} \right) \frac{1}{(1-q)^k} \\ &= \frac{1-q}{(1-q)^k} \sum_{l=0}^k \binom{k}{l} (-1)^l \frac{1 - (\lambda q^{l+1})^{p^N}}{1 - \lambda q^{l+1}} \\ &= \frac{1}{(1-q)^{k-1}} \sum_{l=0}^k \binom{k}{l} (-1)^l \frac{l+1}{1 - \lambda q^{l+1}}. \end{aligned}$$

Therefore, we obtain the following theorem.

Theorem 1. For $k \in \mathbb{N} \cup \{0\}$ and $\lambda \in T_p$, we have

$$\beta_{k,q}(\lambda) = \frac{1}{(1-q)^{k-1}} \sum_{l=0}^k \binom{k}{l} (-1)^l \frac{l+1}{1 - \lambda q^{l+1}}.$$

Let $F(t, \lambda : q)$ be the generating functions of $\beta_{n,q}(\lambda)$ with

$$F(t, \lambda : q) = \sum_{n=0}^{\infty} \beta_{n,q}(\lambda) \frac{t^n}{n!}.$$

Then we have

$$\begin{aligned}
F(t, \lambda : q) &= \sum_{n=0}^{\infty} \beta_{n,q}(\lambda) \frac{t^n}{n!} = \int_{\mathbb{Z}_p} \lambda^x e^{[x]_q t} d\mu_q(x) \\
&= \sum_{n=0}^{\infty} \int_{\mathbb{Z}_p} \lambda^x [x]_q^n d\mu_q(x) \frac{t^n}{n!} \\
(15) \quad &= \sum_{k=0}^{\infty} \left\{ \frac{1}{(1-q)^{k-1}} \sum_{l=0}^k \binom{k}{l} (-1)^l (l+1) \sum_{m=0}^{\infty} \lambda^m q^{(l+1)m} \right\} \frac{t^k}{k!} \\
&= \sum_{k=0}^{\infty} \frac{1}{(1-q)^{k-1}} \sum_{m=0}^{\infty} \lambda^m \sum_{l=0}^k \binom{k}{l} (-1)^l (l+1) q^{(l+1)m} \frac{t^k}{k!} \\
&= \sum_{k=0}^{\infty} \frac{1}{(1-q)^{k-1}} \sum_{m=0}^{\infty} q^m \lambda^m \sum_{l=0}^k l \binom{k}{l} (-1)^l q^{lm} \frac{t^k}{k!} \\
&\quad + \sum_{k=0}^{\infty} \frac{1}{(1-q)^{k-1}} \sum_{m=0}^{\infty} q^m \lambda^m \sum_{l=0}^k \binom{k}{l} (-1)^l q^{lm} \frac{t^k}{k!}.
\end{aligned}$$

Since $l \binom{k}{l} = k \binom{k-1}{l-1}$, the first term of the last equation in (15) equals

$$\begin{aligned}
&\sum_{m=0}^{\infty} q^m \lambda^m \left\{ \sum_{k=0}^{\infty} \frac{1}{(1-q)^{k-1}} \sum_{l=1}^k \binom{k-1}{l-1} (-1)^l q^{lm} \right\} \frac{t^k}{(k-1)!} \\
(16) \quad &= - \sum_{m=0}^{\infty} q^{2m} \lambda^m \left\{ \sum_{k=1}^{\infty} \frac{1}{(1-q)^{k-1}} \sum_{l=0}^{k-1} \binom{k-1}{l} (-1)^l q^{lm} \right\} \frac{t^k}{(k-1)!} \\
&= -t \sum_{m=0}^{\infty} q^{2m} \lambda^m \sum_{k=0}^{\infty} [m]_q^k \frac{t^k}{k!} = -t \sum_{m=0}^{\infty} q^{2m} \lambda^m e^{[m]_q t}.
\end{aligned}$$

The second term of the last equation in (15) equals

$$\begin{aligned}
&\sum_{k=0}^{\infty} \frac{1}{(1-q)^{k-1}} \sum_{m=0}^{\infty} q^m \lambda^m (1-q^m)^k \frac{t^k}{k!} \\
(17) \quad &= (1-q) \sum_{m=0}^{\infty} q^m \lambda^m \sum_{k=0}^{\infty} [m]_q^k \frac{t^k}{k!} = (1-q) \sum_{m=0}^{\infty} q^m \lambda^m e^{[m]_q t}.
\end{aligned}$$

From (15), (16) and (17), we obtain the following proposition.

Proposition 2. *Let $F(t, \lambda : q) = \sum_{n=0}^{\infty} \beta_{n,q}(\lambda) \frac{t^n}{n!}$. Then we have*

$$F(t, \lambda : q) = -t \sum_{m=0}^{\infty} q^{2m} \lambda^m e^{[m]_q t} + (1-q) \sum_{m=0}^{\infty} q^m \lambda^m e^{[m]_q t}.$$

Since $q^{2m} = q^m \{[m]_q(q-1) + 1\}$, it follows that

$$\begin{aligned}
\beta_{k,q}(\lambda) &= \frac{d^k F_q(t, \lambda : q)}{(dt)^k} \Big|_{t=0} \\
&= -k \sum_{m=0}^{\infty} q^{2m} \lambda^m [m]_q^{k-1} + (1-q) \sum_{m=0}^{\infty} q^m \lambda^m [m]_q^k \\
&= -k(q-1) \sum_{m=0}^{\infty} q^m \lambda^m [m]_q^k - k \sum_{m=0}^{\infty} q^m \lambda^m [m]_q^{k-1} + (1-q) \sum_{m=0}^{\infty} q^m \lambda^m [m]_q^k \\
&= (1-q)(k+1) \sum_{m=0}^{\infty} q^m \lambda^m [m]_q^k - k \sum_{m=0}^{\infty} q^m \lambda^m [m]_q^{k-1}.
\end{aligned}$$

Therefore, we obtain the following theorem.

Theorem 3. For $k \in \mathbb{N} \cup \{0\}$ and $\lambda \in T_p$, we have

$$\beta_{k,q}(\lambda) = (1-q)(k+1) \sum_{m=0}^{\infty} q^m \lambda^m [m]_q^k - k \sum_{m=0}^{\infty} q^m \lambda^m [m]_q^{k-1}.$$

Now we consider another q -extension of λ -Bernoulli numbers as follows.

$$(18) \quad B_{n,q}(\lambda) = \int_{\mathbb{Z}_p} q^{-x} \lambda^x [x]_q^n d\mu_q(x).$$

From (18), we can derive

$$\begin{aligned}
B_{n,q}(\lambda) &= \int_{\mathbb{Z}_p} q^{-x} \lambda^x [x]_q^n d\mu_q(x) \\
&= \frac{1}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} \int_{\mathbb{Z}_p} q^{-x} (-1)^l \lambda^x q^{lx} d\mu_q(x) \\
&= \frac{1}{(1-q)^{n-1}} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{l}{1-\lambda q^l}.
\end{aligned}$$

Thus, we obtain the following theorem.

Theorem 4. For $n \in \mathbb{N} \cup \{0\}$ and $\lambda \in T_p$, we have

$$B_{n,q}(\lambda) = \frac{1}{(1-q)^{n-1}} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{l}{1-\lambda q^l}.$$

Let $F^*(t, \lambda : q)$ be the generating functions of $B_{n,q}(\lambda)$ with

$$F^*(t, \lambda : q) = \sum_{n=0}^{\infty} B_{n,q}(\lambda) \frac{t^n}{n!}.$$

Then we have

$$\begin{aligned}
F^*(t, \lambda : q) &= \sum_{n=0}^{\infty} B_{n,q}(\lambda) \frac{t^n}{n!} = \int_{\mathbb{Z}_p} q^{-x} \lambda^x e^{[x]_q t} d\mu_q(x) \\
&= \sum_{n=0}^{\infty} \left\{ \int_{\mathbb{Z}_p} q^{-x} \lambda^x [x]_q^n d\mu_q(x) \right\} \frac{t^n}{n!} \\
&= \sum_{n=0}^{\infty} \left\{ \frac{1}{(1-q)^{n-1}} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{l}{1-\lambda q^l} \right\} \frac{t^n}{n!} \\
&= \sum_{n=0}^{\infty} \left\{ \frac{1}{(1-q)^{n-1}} \sum_{l=0}^n \binom{n}{l} (-1)^l l \sum_{m=0}^{\infty} \lambda^m q^{lm} \right\} \frac{t^n}{n!} \\
&= \sum_{m=0}^{\infty} \lambda^m \left\{ \sum_{n=1}^{\infty} \frac{n}{(1-q)^{n-1}} \sum_{l=1}^n \binom{n-1}{l-1} (-1)^l q^{lm} \right\} \frac{t^n}{n!} \\
&= - \sum_{m=0}^{\infty} \lambda^m q^m \sum_{n=1}^{\infty} \frac{n}{(1-q)^{n-1}} (1-q^m)^{n-1} \frac{t^n}{n!} \\
&= - \sum_{m=0}^{\infty} \lambda^m q^m \sum_{n=0}^{\infty} \frac{(1-q^m)^n}{(1-q)^n} \frac{t^{n+1}}{n!} \\
&= -t \sum_{m=0}^{\infty} \lambda^m q^m e^{[m]_q t}.
\end{aligned}$$

Therefore we obtain the following lemma.

Lemma 5. *Let $F^*(t, \lambda : q) = \sum_{n=0}^{\infty} B_{n,q}(\lambda) \frac{t^n}{n!}$. Then we have*

$$F^*(t, \lambda : q) = -t \sum_{m=0}^{\infty} \lambda^m q^m e^{[m]_q t}.$$

We also have

$$B_{k,q}(\lambda) = \frac{d^k F_q(t, \lambda : q)}{(dt)^k} \Big|_{t=0} = -k \sum_{m=0}^{\infty} q^m \lambda^m [m]_q^{k-1}.$$

Therefore we obtain the following theorem.

Theorem 6. *For $k \in \mathbb{N} \cup \{0\}$ and $\lambda \in T_p$, we have*

$$B_{k,q}(\lambda) = -k \sum_{m=0}^{\infty} q^m \lambda^m [m]_q^{k-1}.$$

3. q -EXTENSION OF λ -EULER NUMBERS AND POLYNOMIALS

In this section, we assume that p is an odd prime number and $\lambda \in \mathbb{C}_p$ with $|1 - \lambda|_p < 1$. By using the fermionic p -adic q -integral on $\mathbb{Z}_p$, we consider the q -extensions of λ -Euler numbers as follows.

For $n \in \mathbb{N} \cup \{0\}$, we define the q -extension of λ -Euler numbers as

$$(19) \quad E_{n,q}(\lambda) = \int_{\mathbb{Z}_p} q^{-x} \lambda^x [x]_q^n d\mu_{-q}(x).$$

From (19), we note that

$$\begin{aligned}
E_{n,q}(\lambda) &= \int_{\mathbb{Z}_p} q^{-x} \lambda^x [x]_q^n d\mu_{-q}(x) \\
&= \lim_{N \rightarrow \infty} \frac{1+q}{1+q^{p^N}} \sum_{x=0}^{p^N-1} (-1)^x [x]_q^n \lambda^x \\
&= \frac{[2]_q}{2} \frac{1}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \lim_{N \rightarrow \infty} \frac{1+q^{p^N} \lambda^{p^N}}{1+q^l \lambda} \\
&= \frac{[2]_q}{2} \frac{1}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{2}{1+q^l \lambda} \\
&= \frac{[2]_q}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{1}{1+q^l \lambda}.
\end{aligned}$$

Therefore we obtain the following theorem.

Theorem 7. *For $n \in \mathbb{N} \cup \{0\}$, we have*

$$E_{n,q}(\lambda) = \frac{[2]_q}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{1}{1+q^l \lambda}.$$

Let $g(t, \lambda : q)$ be the generating function of $E_{n,q}(\lambda)$ with

$$g(t, \lambda : q) = \sum_{n=0}^{\infty} E_{n,q}(\lambda) \frac{t^n}{n!}.$$

Then we have

$$\begin{aligned}
g(t, \lambda : q) &= \sum_{n=0}^{\infty} E_{n,q}(\lambda) \frac{t^n}{n!} = \int_{\mathbb{Z}_p} q^{-x} \lambda^x e^{[x]_q t} d\mu_{-q}(x) \\
&= \sum_{n=0}^{\infty} \left\{ \int_{\mathbb{Z}_p} q^{-x} \lambda^x [x]_q^n d\mu_{-q}(x) \right\} \frac{t^n}{n!} \\
&= [2]_q \sum_{n=0}^{\infty} \left\{ \frac{1}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{1}{1+\lambda q^l} \right\} \frac{t^n}{n!} \\
&= [2]_q \sum_{n=0}^{\infty} \frac{1}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \left\{ \sum_{m=0}^{\infty} (-1)^m \lambda^m q^{lm} \right\} \frac{t^n}{n!} \\
&= [2]_q \sum_{m=0}^{\infty} (-1)^m \lambda^m \sum_{n=0}^{\infty} \frac{1}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lm} \frac{t^n}{n!} \\
&= [2]_q \sum_{m=0}^{\infty} (-1)^m \lambda^m \sum_{n=0}^{\infty} [m]_q^n \frac{t^n}{n!} \\
&= [2]_q \sum_{m=0}^{\infty} (-1)^m \lambda^m e^{[m]_q t}.
\end{aligned}$$

Thus, we have the following lemma.

Lemma 8. Let $g(t, \lambda : q) = \sum_{n=0}^{\infty} E_{n,q}(\lambda) \frac{t^n}{n!}$. Then we have

$$(20) \quad g(t, \lambda : q) = [2]_q \sum_{m=0}^{\infty} (-1)^m \lambda^m e^{[m]_q t}.$$

By (20), we can also consider the λ - q -Genocchi numbers as follows.

$$(21) \quad t \int_{\mathbb{Z}_p} q^{-x} \lambda^x e^{[x]_q t} d\mu_{-q}(x) = [2]_q t \sum_{m=0}^{\infty} (-1)^m \lambda^m e^{[m]_q t} = \sum_{n=0}^{\infty} G_{n,q}(\lambda) \frac{t^n}{n!}.$$

From (21), we note that $G_{0,q}(\lambda) = 0$ and

$$\int_{\mathbb{Z}_p} q^{-x} \lambda^x [x]_q^n d\mu_{-q}(x) = \frac{G_{n+1,q}(\lambda)}{n+1}.$$

Thus, we see that

$$E_{n,q}(\lambda) = \int_{\mathbb{Z}_p} q^{-x} \lambda^x [x]_q^n d\mu_{-q}(x) = \frac{G_{n+1,q}(\lambda)}{n+1}.$$

Hence

$$G_{n,q}(\lambda) = [2]_q \frac{n}{(1-q)^{n-1}} \sum_{l=0}^{n-1} \binom{n-1}{l} (-1)^l \frac{1}{1+q^l \lambda},$$

where $n = 1, 2, 3, \dots$. Indeed,

$$\begin{aligned} G_{1,q}(\lambda) &= \frac{[2]_q}{1+\lambda}, \\ G_{2,q}(\lambda) &= \frac{2[2]_q}{1-q} \sum_{l=0}^1 \binom{1}{l} (-1)^l \frac{1}{1+q^l \lambda} = \frac{[2]_q}{1-q} \left(\frac{2}{1+\lambda} - \frac{2}{1+q\lambda} \right) \\ &= -2[2]_q \left(\frac{\lambda}{(1+\lambda)(1+q\lambda)} \right). \end{aligned}$$

Now, we consider the q -extension of λ -Euler polynomials as follows.

$$(22) \quad E_{n,q}(\lambda, x) = \int_{\mathbb{Z}_p} q^{-y} \lambda^y [x+y]_q^n d\mu_{-q}(y).$$

From (22), we can easily derive

$$E_{n,q}(\lambda, x) = \frac{[2]_q}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} \frac{1}{1+q^l \lambda}.$$

Let $g(x, \lambda : q) = \sum_{n=0}^{\infty} E_{n,q}(\lambda, x) \frac{t^n}{n!}$. Then we have

$$\begin{aligned} g(x, \lambda : q) &= \sum_{n=0}^{\infty} E_{n,q}(\lambda, x) \frac{t^n}{n!} = \int_{\mathbb{Z}_p} q^{-y} \lambda^y e^{[x+y]_q t} d\mu_{-q}(y) \\ &= \sum_{n=0}^{\infty} \frac{[2]_q}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} \left(\sum_{m=0}^{\infty} (-1)^m q^{ml} \lambda^m \right) \frac{t^n}{n!} \\ &= [2]_q \sum_{m=0}^{\infty} (-1)^m \lambda^m e^{[m+x]_q t}. \end{aligned}$$

It follows that

$$E_{n,q}(\lambda, x) = \frac{d^n(g(x, \lambda : g))}{(dt)^n} \Big|_{t=0} = [2]_q \sum_{m=0}^{\infty} (-1)^m \lambda^m [m+x]_q^n.$$

Then we obtain the following theorem.

Theorem 9. *For $n \in \mathbb{N} \cup \{0\}$, we have*

$$E_{n,q}(\lambda, x) = [2]_q \sum_{m=0}^{\infty} (-1)^m \lambda^m [m+x]_q^n.$$

By the same method, we consider the λ - q -Genocchi polynomials as follows.

$$\begin{aligned} (23) \quad t \int_{\mathbb{Z}_p} q^{-x} \lambda^x e^{[x+y]_q t} d\mu_{-q}(x) &= [2]_q t \sum_{m=0}^{\infty} (-1)^m \lambda^m e^{[m+x]_q t} \\ &= \sum_{n=0}^{\infty} G_{n,q}(\lambda, x) \frac{t^n}{n!}. \end{aligned}$$

By (23), we see

$$(24) \quad E_{n,q}(\lambda, x) = \int_{\mathbb{Z}_p} q^{-y} \lambda^y [x+y]_q^n d\mu_{-q}(y) = \frac{G_{n+1,q}(\lambda, x)}{n+1}$$

and $G_{0,q}(\lambda, x) = 0$. From the definition of λ - q -Euler polynomials, we derive

$$\begin{aligned} (25) \quad E_{n,q}(\lambda, x) &= \int_{\mathbb{Z}_p} q^{-y} \lambda^y [x+y]_q^n d\mu_{-q}(y) \\ &= \sum_{l=0}^n \binom{n}{l} [x]_q^{n-l} q^{lx} \int_{\mathbb{Z}_p} q^{-y} \lambda^y [y]_q^l d\mu_{-q}(y) \\ &= \frac{[2]_q}{[2]_{q^d}} [d]_q^n \sum_{a=0}^{d-1} (-1)^a \lambda^a \int_{\mathbb{Z}_p} q^{-dy} \lambda^{dy} \left[\frac{x+a}{d} + y \right]_{q^d}^n d\mu_{-q}(y,) \end{aligned}$$

for $d \in \mathbb{N}$ with $d \equiv 1 \pmod{2}$. By (25), we see that

$$\begin{aligned} E_{n,q}(\lambda, x) &= \sum_{l=0}^n \binom{n}{l} [x]_q^{n-l} q^{lx} E_{l,q}(\lambda, x) \\ &= \frac{[2]_q}{[2]_{q^d}} [d]_q^n \sum_{a=0}^{d-1} (-1)^a \lambda^a E_{n,q^d}(\lambda^d, \frac{x+a}{d}). \end{aligned}$$

It is easy to see that

$$qI_{-q}(f_1) + I_{-q}(f) = [2]_q f(0),$$

where $f_1(x) = f(x+1)$. Thus, we have

$$q \int_{\mathbb{Z}_p} q^{-y-1} \lambda^{y+1} [x+1+y]_q^n d\mu_{-q}(y) + \int_{\mathbb{Z}_p} q^{-y} \lambda^y [x+y]_q^n d\mu_{-q}(y) = [2]_q [x]_q^n.$$

Therefore, we obtain the following theorem.

Theorem 10. *For $n \in \mathbb{N} \cup \{0\}$, we have*

$$\lambda E_{n,q}(\lambda, x+1) + E_{n,q}(\lambda, x) = [2]_q [x]_q^n.$$

By Theorem 10 and (24), we have the following result.

Corollary 11. *For $n \in \mathbb{N} \cup \{0\}$, we have*

$$\lambda G_{n,q}(\lambda, x+1) + G_{n,q}(\lambda, x) = [2]_q n [x]_q^{n-1}.$$

It is easy to see that

$$\begin{aligned} \frac{\partial}{\partial x} [x+y]_q^n &= n [x+y]_q^{n-1} \frac{\log q}{q-1} q^{x+y} \\ &= n \log q [x+y]_q^{n-1} + \frac{\log q}{q-1} n [x+y]_q^{n-1}. \end{aligned}$$

From (22), we note that

$$(26) \quad \frac{\partial}{\partial x} E_{n,q}(\lambda, x) = \frac{\partial}{\partial x} \int_{\mathbb{Z}_p} q^{-y} \lambda^y [x+y]_q^n d\mu_{-q}(y).$$

The right side of (26) equals

$$\begin{aligned} n \log q \int_{\mathbb{Z}_p} q^{-y} \lambda^y [x+y]_q^n d\mu_{-q}(y) + \frac{\log q}{q-1} n \int_{\mathbb{Z}_p} q^{-y} \lambda^y [x+y]_q^n d\mu_{-q}(y) \\ = n \log q E_{n,q}(\lambda, x) + \frac{\log q}{q-1} n E_{n-1,q}(\lambda, x). \end{aligned}$$

Therefore, we obtain the following lemma.

Lemma 12. *For $n \in \mathbb{N}$, we have*

$$\frac{\partial}{\partial x} E_{n,q}(\lambda, x) = n \log q E_{n,q}(\lambda, x) + \frac{\log q}{q-1} n E_{n-1,q}(\lambda, x).$$

Remark 1. *Note that*

$$\begin{aligned} \frac{\partial}{\partial x} G_{n,q}(\lambda, x) &= n E_{n-1,q}(\lambda, x) \\ &= \frac{n}{(1-q)^{n-1}} [2]_q \sum_{l=0}^{n-1} \binom{n-1}{l} (-1)^l q^{lx} \frac{1}{1+q^l \lambda}. \end{aligned}$$

Remark 2. *Note that*

$$\begin{aligned} E_{n,q}(\lambda, dx) &= \int_{\mathbb{Z}_p} q^{-y} \lambda^y [dx+y]_q^n d\mu_{-q}(y) \\ &= [d]_q^n \frac{[2]_q}{[2]_{q^d}} \sum_{a=0}^{n-1} (-1)^a \lambda^a \int_{\mathbb{Z}_p} [x + \frac{a}{d} + y]_{q^d}^n \lambda^{dy} q^{-dy} d\mu_{-q^d}(y) \\ &= [d]_q^n \frac{[2]_q}{[2]_{q^d}} \sum_{a=0}^{n-1} (-1)^a \lambda^a E_{n,q^d}(\lambda^d, x + \frac{a}{d}), \end{aligned}$$

for $d \in \mathbb{N}$ with $d \equiv 1 \pmod{2}$.

For $n \in \mathbb{N}$, it is known that

$$(27) \quad q^n I_{-q}(f_n) = (-1)^n I_{-q}(f) + [2]_q \sum_{l=0}^{n-1} (-1)^{n-1-l} q^l f(l), \quad (\text{see [7]}),$$

where $f_n(x) = f(x+n)$. By (27), we obtain the following lemma.

Lemma 13. *For $n \in \mathbb{N}$, we have*

$$q^n I_{-q}(f_n) + (-1)^{n-1} I_{-q}(f) = [2]_q \sum_{l=0}^{n-1} (-1)^{n-1-l} q^l f(l).$$

For $n \equiv 1 \pmod{2}$, we also have

$$q^n I_{-q}(f_n) + I_{-q}(f) = [2]_q \sum_{l=0}^{n-1} (-1)^l q^l f(l).$$

If we take $f(x) = \lambda^x q^{-x} [x]_q^m$ with $m \in \mathbb{N} \cup \{0\}$, then we see that

$$q^n \int_{\mathbb{Z}_p} q^{-x-n} \lambda^{x+n} [x+n]_q^m d\mu_{-q}(x) + \int_{\mathbb{Z}_p} q^{-x} \lambda^x [x]_q^m d\mu_{-q}(x) = [2]_q \sum_{l=0}^{n-1} (-1)^l \lambda^l [l]_q^m.$$

Thus we have

$$E_{m,q}(\lambda, n) + E_{m,q}(\lambda) = [2]_q \sum_{l=0}^{n-1} (-1)^l \lambda^l [l]_q^m.$$

For $m \equiv 1 \pmod{2}$, we note that

$$\begin{aligned} E_{n,q}(\lambda, dx) &= \frac{[2]_q}{[2]_{q^m}} [m]_q^n \sum_{a=0}^{m-1} (-1)^a \lambda^a E_{n,q^m}(\lambda^m, \frac{a}{m}) \\ &= \frac{[2]_q}{[2]_{q^m}} [m]_q^n \sum_{a=0}^n (-1)^a \lambda^a \sum_{l=0}^n \binom{n}{l} \left(\frac{[a]_q}{[m]_q} \right)^{n-l} q^{la} E_{n,q^m}(\lambda^m, \frac{a}{m}) \\ &= \frac{[2]_q}{[2]_{q^m}} \sum_{l=0}^n \binom{n}{l} [m]_q^l E_{l,q^m}(\lambda^m) \sum_{a=0}^{m-1} (-1)^a \lambda^a q^{la} [a]_q^{n-l}. \end{aligned}$$

Remark 3. *Note that*

$$\frac{G_{m+1,q}(\lambda, n)}{m+1} + \frac{G_{m+1,q}(\lambda)}{m+1} = [2]_q \sum_{l=0}^{n-1} (-1)^l \lambda^l [l]_q^m.$$

Now we can also consider the following DC type λ - q -Euler numbers and polynomials. For $\lambda \in \mathbb{C}_p$ with $|1 - \lambda|_p < 1$, we define the DC type λ - q -Euler numbers as

$$\begin{aligned} E_{n,q}^*(\lambda) &= \int_{\mathbb{Z}_p} \lambda^x [x]_q^n d\mu_{-q}(x) \\ &= \frac{[2]_q}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l \frac{1}{1 + q^{l+1} \lambda} \\ &= [2]_q \sum_{m=0}^{\infty} (-1)^m \lambda^m q^m [m]_q^n. \end{aligned}$$

Let $g^*(t, \lambda : q) = \sum_{n=0}^{\infty} E_{n,q}^*(\lambda) \frac{t^n}{n!}$. Then we see that

$$g^*(t, \lambda : q) = \int_{\mathbb{Z}_p} \lambda^x e^{[x]_q t} d\mu_{-q}(x) = [2]_q \sum_{m=0}^{\infty} (-1)^m \lambda^m q^m e^{[m]_q t}.$$

The DC type λ - q -Euler polynomials are also defined as

$$\begin{aligned} E_{n,q}^*(\lambda, x) &= \int_{\mathbb{Z}_p} \lambda^y [x+y]_q^n d\mu_{-q}(y) \\ &= \frac{[2]_q}{(1-q)^n} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} \frac{1}{1+q^{l+1}\lambda} \\ &= [2]_q \sum_{m=0}^{\infty} (-1)^m \lambda^m q^m [m+x]_q^n. \end{aligned}$$

Thus we can give the generating function of the DC type λ - q -Euler polynomials as follows.

$$\sum_{n=0}^{\infty} E_{n,q}^*(\lambda, x) \frac{t^n}{n!} = \int_{\mathbb{Z}_p} \lambda^y e^{[x+y]_q t} d\mu_{-q}(y) = [2]_q \sum_{m=0}^{\infty} (-1)^m \lambda^m q^m e^{[m+x]_q t}.$$

4. FURTHER REMARKS AND OBSERVATION FOR THE q -EXTENSION LERCH ZETA FUNCTION

In this section, we assume that $q \in \mathbb{C}$ with $|q| < 1$. It is well-known that Lerch type zeta function is defined as

$$\zeta(x, s, a) = \sum_{n=0}^{\infty} \frac{x^n}{(n+a)^s},$$

where $a \in \mathbb{C}$ with $a \neq 0, -1, -2, \dots$, and $s \in \mathbb{C}$ when $|x| < 1$, $Re(s) > 1$ when $|x| = 1$, and Hurwitz zeta function is defined as

$$\zeta(s, a) = \sum_{n=0}^{\infty} \frac{1}{(n+a)^s},$$

where $Re(s) > 1$ and $a \neq 0, -1, -2, \dots$. The Lerch zeta function is known that

$$\zeta(s, \eta) = \sum_{n=0}^{\infty} \frac{e^{2\pi i n \eta}}{n^s} = e^{2\pi i \eta} \zeta(e^{2\pi i \eta}, s, 1),$$

where $\eta \in \mathbb{R}$ and $Re(s) > 1$.

Now we consider the first kind of the q -extension of Lerch type zeta function as follows.

$$(28) \quad \zeta_q(\lambda, s) = (1-q) \frac{2-s}{s-1} \sum_{m=1}^{\infty} \frac{q^m \lambda^m}{[m]_q^{s-1}} + \sum_{m=1}^{\infty} \frac{q^m \lambda^m}{[m]_q^s},$$

where $q \in \mathbb{C}$ with $|q| < 1$, and $\lambda \in \mathbb{C}$ with $\lambda = e^{2\pi i/f}$ ($f \in \mathbb{N}$).

By Theorem 3, we see that

$$(29) \quad -\frac{\beta_{k,q}(\lambda)}{k} = (q-1) \frac{k+1}{k} \sum_{m=1}^{\infty} q^m \lambda^m [m]_q^k + \sum_{m=1}^{\infty} q^m \lambda^m [m]_q^{k-1},$$

for $k \in \mathbb{N}$.

By (28) and (29), we obtain the following theorem.

Theorem 14. For $k \in \mathbb{N}$, we have

$$\zeta_q(\lambda, 1 - k) = -\frac{\beta_{k,q}(\lambda)}{k}.$$

Now, we define the second of the q -extension of Lerch zeta function as follows. For $s \in \mathbb{C}$ and $\lambda = e^{2\pi i/f}$ ($f \in \mathbb{N}$), define

$$(30) \quad \zeta_q^*(\lambda, s) = \sum_{m=1}^{\infty} \frac{q^m \lambda^m}{[m]_q^s}.$$

By Theorem 6, we easily see that

$$(31) \quad -\frac{\beta_{k,q}(\lambda)}{k} = \sum_{m=1}^{\infty} q^m \lambda^m [m]_q^{k-1}.$$

By (30) and (31), we obtain the following theorem.

Theorem 15. For $k \in \mathbb{N}$, we have

$$\zeta_q^*(\lambda, 1 - k) = -\frac{\beta_{k,q}(\lambda)}{k}.$$

Remark 4. The extension of Hurwitz's type q -Euler zeta function is defined as

$$\zeta_{q,E}(\lambda, s) = [2]_q \sum_{m=0}^{\infty} \frac{(-1)^m \lambda^m}{[m+x]_q^s},$$

where $s \in \mathbb{C}$, $\lambda \in \mathbb{C}$ with $\lambda = e^{2\pi i/f}$ ($f \in \mathbb{N}$). Then we have

$$\zeta_q(\lambda, 1 - k) = E_{k,q}(\lambda, x), \quad (k \in \mathbb{N}).$$

REFERENCES

- [1] I. N. Cangul, H. Ozden, Y. Simsek, *Generating functions of the (h, q) extension of twisted Euler polynomials and numbers*, Acta Math. Hungar. **120** (2008), no. 3, 281–299.
- [2] M. Cenkci, V. Kurt, *Congruences for generalized q -Bernoulli polynomials*, J. Inequal. Application 2008, Art. ID 270713, 19 pp.
- [3] M. Cenkci, Y. Simsek, V. Kurt, *Further remarks on multiple p -adic q - L -function of two variables*, Adv. Stud. Contemp. Math. (Kyungshang) **14** (2007), no. 1, 49–68.
- [4] M. Cenkci, Y. Simsek, V. Kurt, *Multiple two-variable p -adic q - L -function and its behavior at $s = 0$* , Russian Journal of Mathematical Physics **15** (2008) no. 4, 447–459.
- [5] T. Kim, *An analogue of Bernoulli numbers and their congruences*, Rep. Fac. Sci. Engrg. Saga Univ. Math. **22** (1994), no. 2, 21–26.
- [6] T. Kim, *On a q -analogue of the p -adic log gamma functions and related integrals*, J. Number Theory **76** (1999), no. 2, 320–329.
- [7] T. Kim, *q -Euler numbers and polynomials associated with p -adic q -integrals*, J. Nonlinear Math. Phys. **14** (2007), no. 1, 15–27.
- [8] T. Kim, *q -Bernoulli numbers and polynomials associated with Gaussian binomial coefficients*, Russ. J. Math. Phys. **15** (2008), no. 1, 51–57.
- [9] T. Kim, *On the multiple q -Genocchi and Euler numbers*, Russian Journal of Mathematical Physics **15** (2008) no.4, 481–486.
- [10] T. Kim, S.-H. Rim, Y. Simsek, D. Kim, *On the analogs of Bernoulli and Euler numbers, related identities and zeta and L -functions*, J. Korean Math. Soc. **45** (2008), no. 2, 435–453.
- [11] Y.-H. Kim, W. Kim, and L.-C. Jang, *On the q -extension of Apostol-Euler numbers and polynomials*, Abstract and Applied Analysis (2008)
<http://www.hindawi.com/journals/aaa/aip.296159.html>.

- [12] H. Ozden, Y. Simsek, *A new extension of q -Euler numbers and polynomials related to their interpolation functions*, Appl. Math. Lett. **21** (2008), no. 9, 934–939.
- [13] H. Ozden, Y. Simsek, *Interpolation function of the (h, q) -extension of twisted Euler numbers*, Comput. Math. Appl. 56 (2008), no. 4, 898–908.
- [14] H. Ozden, I. N. Cangul, Y. Simsek, *Multivariate interpolation functions of higher-order q -Euler numbers and their applications*, Abstr. Appl. Anal. 2008, Art. ID 390857, 16 pp.
- [15] K. H. Park, Y.-H. Kim, *On some arithmetical properties of the Genocchi numbers and polynomials*, Advances in Difference Equations (2008)
<http://www.hindawi.com/journals/ade/aip.195049.html>.
- [16] Y. Simsek, *q -analogue of twisted l -series and q -twisted Euler numbers*, J. Number Theory **110** (2005), no. 2, 267–278.
- [17] Y. Simsek, *On p -adic twisted q - L -functions related to generalized twisted Bernoulli numbers*, Russ. J. Math. Phys. **13** (2006), no. 3, 340–348.
- [18] Y. Simsek, *Twisted (h, q) -Bernoulli numbers and polynomials related to twisted (h, q) -zeta function and L -function*, J. Math. Anal. Appl. **324** (2006), no. 2, 790–804.
- [19] Y. Simsek, *Generating functions of the twisted Bernoulli numbers and polynomials associated with their interpolation functions*, Adv. Stud. Contemp. Math. (Kyungshang) **16** (2008), no. 2, 251–278.

TAEKYUN KIM. DIVISION OF GENERAL EDUCATION-MATHEMATICS,, KWANGWOON UNIVERSITY,
SEOUL 139-701, REPUBLIC OF KOREA,
E-mail address: `tkkim@kw.ac.kr`

YOUNG-HEE KIM. DIVISION OF GENERAL EDUCATION-MATHEMATICS,, KWANGWOON UNIVERSITY,
SEOUL 139-701, REPUBLIC OF KOREA,
E-mail address: `yhkim@kw.ac.kr`

KYUNG-WON HWANG. DEPARTMENT OF GENERAL EDUCATION,, KOOKMIN UNIVERSITY, 861-1
SEONGBUKGU SEOUL 136-702, REPUBLIC OF KOREA,
E-mail address: `khwang7@kookmin.ac.kr`